

TRIPLE CORRELATIONS OF MULTIPLICATIVE FUNCTIONS

PRANENDU DARBAR

ABSTRACT. In this paper, we find asymptotic formula for the following sum with explicit error term:

$$M_x(g_1, g_2, g_3) = \frac{1}{x} \sum_{n \leq x} g_1(F_1(n))g_2(F_2(n))g_3(F_3(n)),$$

where $F_1(x), F_2(x)$ and $F_3(x)$ are polynomials with integer coefficients and g_1, g_2, g_3 are multiplicative functions with modulus less than or equal to 1.

Moreover, under some assumption on g_1, g_2 , we prove that as $x \rightarrow \infty$,

$$\frac{1}{x} \sum_{n \leq x} g_1(n+3)g_2(n+2)\mu(n+1) = o(1)$$

and assuming 2-point Chowla type conjecture we show that as $x \rightarrow \infty$,

$$\frac{1}{x} \sum_{n \leq x} g_1(n+3)\mu(n+2)\mu(n+1) = o(1).$$

1. INTRODUCTION

Let $g_j : \mathbb{N} \rightarrow \mathbb{C}$ be multiplicative functions such that $|g_j(n)| \leq 1$ for all n and $j = 1, 2, 3$. Let $F_1(x), F_2(x), F_3(x)$ are polynomials with integer coefficients.

Consider the following triple correlation function:

$$(1) \quad M_x(g_1, g_2, g_3) = \frac{1}{x} \sum_{n \leq x} g_1(F_1(n))g_2(F_2(n))g_3(F_3(n)),$$

In [4], Katái studied the asymptotic behaviour of the above sum (1) when $F_j(x), j = 1, 2, 3$ are special polynomials and some assumptions on $g_j, j = 1, 2, 3$, but did not provide error term. In [13], Stepanauskas studied the asymptotic formula for sum (1) with explicit error term when $F_j(x), j = 1, 2, 3$ are linear polynomials and g_1, g_2, g_3 are close to 1 (see definition 1). Recently, Klurman [15] studied the double correlation function (i.e. the sum (1) with $g_3 = 1$).

Estimations of (1) can be used to get information on the behaviour of the limit distribution of the sum

$$(2) \quad f_1(F_1(n)) + f_2(F_2(n)) + f_3(F_3(n)),$$

where f_1, f_2 and f_3 are real-valued additive functions.

From onwards, let $F(n); F_1(n), F_2(n), F_3(n)$ be positive integer-valued polynomials with integer coefficients and these are not divisible by the square of any irreducible polynomial. Also suppose that $F_j(n), F_k(n)$ are relatively primes for $j \neq k$ and for all n . Let v and v_j denote the degree of the polynomials $F(n)$ and $F_j(n)$ respectively.

In this paper, we will investigate the following sums with various assumptions on $g_j, j = 1, 2, 3$:

$$(3) \quad M'_x(g_1, g_2, g_3) = \frac{1}{x} \sum_{n \leq x} g_1(n+3)g_2(n+2)g_3(n+1)$$

and asymptotic formula of the following triple correlation function with explicit error term which is a improvement of a theorem of Katái ([4], Theorem 5) with respect to explicit error term:

$$(4) \quad M''_x(g_1, g_2, g_3) = \frac{1}{x} \sum_{n \leq x} g_1(F_1(n))g_2(F_2(n))g_3(F_3(n)),$$

where $F_1(x), F_2(x)$ and $F_3(x)$ are polynomials as above of degree ≥ 2 .

Definition 1. A multiplicative function g_j is said to be close to 1 if

$$(5) \quad \sum_p \frac{g_j(p) - 1}{p} < \infty.$$

Definition 2. A multiplicative function g_j is called good function if there exists a $\kappa \in \mathbb{C}$ such that for each $u > 0$

$$(6) \quad \sum_{p \leq x} |g_j(p) - \kappa| \ll \frac{x}{(\log x)^u}.$$

Outline. In section 5, we formulate the sums (3) and (4) in terms of main term and explicit error term.

For the asymptotic behaviour of the sum (3) it seems to be very difficult if we take either g_1, g_2 , and g_3 are equal to Mobius function or any two of them are equal to Mobius function and other one is close to some fixed complex number.

In section 6, firstly we investigate the asymptotic behaviour of the sum (3) when g_1, g_2 are close to 1 and g_3 is the Mobius function. Secondly, we investigate the sum (3) in terms of main term and explicit error term when g_3 is a good function.

In section 7, under some assumption we investigate the asymptotic behaviour of the sum (3) when g_1 is close to 1 and g_2, g_3 are Mobius function.

In section 8, we formulate some applications the above sums (3) and (4).

Acknowledgement. I would like to express my appreciation to my thesis supervisor Anirban Mukhopadhyay for his valuable and constructive suggestions during the planning and development of this paper. I thank Gediminas Stepanauskas for providing me with preprint of [12],[13],[14].

2. Notations

Throughout the paper p and q denote primes; j, k, l, m and n are natural numbers; $c, c', c_1, c_2, \dots$ are absolute constants; $c_F, c_{F_1}, c_{F_2}, c_{F_3}$ are constants depending on F, F_1, F_2, F_3 and $\wp$ is the set of all primes.

In section 3, we use the following notations for Theorem 1 :

$$S(r, x, g_j) = \sum_{r < p \leq x+4-j} \frac{|g_j(p) - 1|^2}{p}, \quad S(r, x) = \sum_{j=1}^3 S(r, x, g_j), \quad r > 0,$$

$$P(x) = \prod_{p \leq x} w_p \quad \text{and} \quad P(r, x) = \prod_{r < p \leq x} w_p,$$

where

$$(7) \quad w_p = \sum_{\substack{m_1=0 \\ (p^{m_1}, 3)=1}}^{\infty} \sum_{\substack{m_2=0 \\ (p^{m_2}, 2)=1 \\ (p^{m_1}, p^{m_2})=1 \\ (p^{m_2}, p^{m_3})=1 \\ (p^{m_1}, p^{m_3})|2}}^{\infty} \sum_{m_3=0}^{\infty} \frac{h_1(p^{m_1})h_2(p^{m_2})h_3(p^{m_3})}{[p^{m_1}, p^{m_2}, p^{m_3}]}.$$

We use the following notations for Theorem 2 :

Let $\varrho(d)$ and $\varrho_j(d)$ denote the number of congruent solutions of the congruences $F(n) \equiv 0 \pmod{d}$ and $F_j(n) \equiv 0 \pmod{d}$ respectively.

Let $\varrho(d_1, d_2, d_3)$ be the number of solutions of the congruence system

$$F_j(n) \equiv 0 \pmod{d_j} \quad j = 1, 2, 3.$$

Let D_γ denote the set of those tuples $\{d_1, d_2, d_3\}$ of natural numbers, for which all the prime factors of d_i do not exceed γ .

$$S'(r, x) = \sum_{r < p \leq x} \sum_{j=1}^3 \frac{|g_j(p) - 1|^2 \varrho_j(p)}{p}, \quad T(x) = \sum_{j=1}^3 \sum_{x < p \leq F_j(x)} \frac{|g_j(p) - 1|^2 \varrho_j(p)}{p},$$

$$C(r) = \sum_{j=1}^3 \sum_{m=1}^{v_j-1} \sum_{\substack{p^m \leq F_j(x) \\ p > r}} |g_j(p^m) - 1| \varrho_j(p^m),$$

$$P'(x) = P_1(\gamma)P_2(\gamma, x) \quad \text{and} \quad P'(r, x) = \prod_{r < p \leq x} w'_p,$$

where

$$(8) \quad P_1(\gamma) = \sum_{\{d_1, d_2, d_3\} \in D_\gamma} \frac{h_1(d_1)h_2(d_2)h_3(d_3)}{[d_1, d_2, d_3]} \varrho(d_1, d_2, d_3),$$

$$(9) \quad P_2(\gamma, x) = \prod_{\gamma < p \leq x} \left(1 + \sum_{m=1}^{\infty} \sum_{j=1}^3 \frac{h_j(p^m) \varrho_j(p^m)}{p^m} \right) := \prod_{\gamma < p \leq x} w'_p.$$

For Theorem 3 and 4 we need the following notations:

$$\theta_\tau(n) = \prod_{p|n} \left(1 + \sum_{m=1}^{\infty} \frac{g_3(p^m)}{p^{m(1+i\tau)}} \right)^{-1}, \quad \tau \in \mathbb{R},$$

$$Q(r) = \prod_{p \leq r} \left(1 - \frac{2\theta_\tau(p)}{p-1} + \theta_\tau(p) \sum_{m=1}^{\infty} \frac{g_1(p^m) + g_2(p^m)}{p^m} \right),$$

$$P''(r, x) = \prod_{r < p \leq x} \left(1 - \frac{1}{p} \right) \left(1 + \sum_{m=1}^{\infty} \frac{g_1(p^m) + g_2(p^m)}{p^m} \right),$$

$$S''(r, x) = \sum_{j=1}^2 \sum_{r < p \leq x+4-j} \frac{|g_j(p) - 1|^2}{p} \quad \text{and} \quad M_x(g_3) = \frac{1}{x} \sum_{n \leq x} g_3(n).$$

3. Statements of Theorems

We begin with the asymptotic formula for the sum (3) with explicit error term which is a special case of a theorem of Stepanauskas([13]).

Theorem 1. *Let g_1, g_2 and g_3 be multiplicative functions with modulus less than or equal to 1. Then there exists a positive absolute constant c such that for all $x \geq r \geq 2$ and for all $\frac{2}{3} < \alpha < 1$, we have*

$$M'_x(g_1, g_2, g_3) = P(x) + O\left(x^{2-3\alpha} \exp\left(c \frac{r^\alpha}{\log r}\right) + (S(r, x))^{\frac{1}{2}} + (r \log r)^{-\frac{1}{2}}\right).$$

The aim of this paper is to prove the following statements:

Theorem 2. *Let $F_j(x), j = 1, 2, 3$ be polynomials as above of degree greater than or equal to 2. Let g_1, g_2 and g_3 be multiplicative functions whose modulus do not exceed 1. Then there exists a positive absolute constant c and a natural number γ such that for all $x \geq r \geq \gamma$ and for all $1 - \frac{1}{v_1+v_2+v_3} < \alpha < 1$, we have*

$$\begin{aligned} M''_x(g_1, g_2, g_3) - P'(x) &\ll (F_1(x)F_2(x)F_3(x))^{1-\alpha} \exp\left(c \frac{r^\alpha}{\log r}\right) + (S'(r, x))^{\frac{1}{2}} \\ &+ (T(x))^{\frac{1}{2}} + (r \log r)^{-\frac{1}{2}} + \frac{1}{x}C(r) + \frac{1}{\log x}. \end{aligned}$$

Theorem 3. *Let g_1 and g_2 be multiplicative functions which do not exceed 1 and*

$$(10) \quad \sum_p \sum_{i=1}^2 \frac{|g_i(p) - 1|^2}{p} < \infty.$$

Then as $x \rightarrow \infty$,

$$M'_x(g_1, g_2, \mu) = \frac{1}{x} \sum_{n \leq x} g_1(n+3)g_2(n+2)\mu(n+1) = o(1).$$

Theorem 4. *Let g_1, g_2 and g_3 be multiplicative functions whose modulus do not exceed 1 and g_3 is a good function.*

Assume further that there exist a positive constant c_1 such that

$$(11) \quad \left| 1 + \sum_{k=1}^{\infty} \frac{g_3(2^k)}{2^{k(1+i\xi)}} \right| \geq c_1$$

for $\xi = 0$, if g_3 is real valued and for all $\xi \in \mathbb{R}$, if g_3 is not real valued. Then there exists a positive absolute constant c and a real $\tau, |\tau| \leq (\log x)^{1/19}$, such that for all $x \geq r \geq 2$ and for all $\frac{1}{2} < \alpha < \frac{5}{9}$, we have

$$\begin{aligned} M'_x(g_1, g_2, g_3) - M_x(g_3)P''(r, x)Q(r) &\ll x^{1-2\alpha} \exp\left(c \frac{r^\alpha}{\log r}\right) + \frac{(\log r)^c}{(\log x)^c} \\ &+ \frac{\exp(c(\log \log r)^2)}{(\log x)^{1/19}} + (S''(r, x))^{\frac{1}{2}} + (r \log r)^{-\frac{1}{2}}. \end{aligned}$$

For real-valued g_3 we may set $\tau = 0$.

Assumption 1 (2-point Chowla type conjecture). *For every given $A > 0$,*

$$\sum_{n \leq x} \mu(n+2)\mu(n+1) \exp(2\pi i n \alpha) = O\left(\frac{x}{(\log x)^A}\right)$$

holds uniformly for all real α .

Theorem 5. *Let g_1 be a multiplicative function such that $|g_1(n)| \leq 1$ for all n and*

$$(12) \quad \sum_p \frac{|g_1(p) - 1|^2}{p} < \infty.$$

Suppose that Assumption 1 holds, then as $x \rightarrow \infty$,

$$M'_x(g_1, \mu, \mu) = \frac{1}{x} \sum_{n \leq x} g_1(n+3)\mu(n+2)\mu(n+1) = o(1).$$

4. Applications

In this section we state several applications of our results.

Application 1 is a linear version of a theorem of Katái ([4], Theorem 1).

Application 1. *Let g_1, g_2, g_3 be multiplicative function such that for all $j = 1, 2, 3$, $|g_j| \leq 1$ and g_1, g_2, g_3 are close to 1. Then as $x \rightarrow \infty$ we have,*

$$M'_x(g_1, g_2, g_3) = \prod_p w_p + o(1),$$

where w_p is defined in (7).

Application 2 is a polynomial version with degree of polynomial greater than or equal to 2 of a theorem of Katái ([4], Theorem 5).

Application 2. *Let $F_j(n)$ ($j = 1, 2, 3$) be as above of degree $v_j \geq 2$. Let g_j ($j = 1, 2, 3$) be as above and*

$$(13) \quad \sum_p \frac{(g_j(p) - 1)\varrho_j(p)}{p} < \infty.$$

Suppose that as $p \rightarrow \infty$,

$$(14) \quad (g_j(p^\alpha) - 1)\varrho_j(p^\alpha) \rightarrow 0,$$

for $\alpha = 1$, when $v_j \geq 2$ and for $\alpha = 1, 2, \dots, v_j - 2$, when $v_j \geq 3$, then we have as $x \rightarrow \infty$,

$$M''_x(g_1, g_2, g_3) \rightarrow P_1(\gamma)P_2(\gamma),$$

where $P_1(\gamma)$ is defined by (8) and

$$(15) \quad P_2(\gamma) := \prod_{p > \gamma} \left(1 + \sum_{m=1}^{\infty} \sum_{i=1}^3 \frac{h_i(p^m)\varrho_i(p^m)}{p^m} \right).$$

Applications 3, 4, 5 and 6 are the direct applications of the Theorems 1, 3, 5 and 2 respectively.

Application 3. *Let $\phi(n) = n \prod_{p|n} \left(1 - \frac{1}{p}\right)$, be Euler's totient function and $\sigma_a(n) =$*

$\sum_{d|n} d^a, a > 0$. Then for $x \geq 2, 0 < A < 1$ we have,

$$\begin{aligned} \frac{1}{x} \sum_{n \leq x} \frac{\sigma_a(n+3)\sigma_a(n+2)\sigma_a(n+1)}{(n+3)^a(n+2)^a(n+1)^a} &= w_2 \prod_{p > 2} \left\{ 1 + \frac{3}{p^{a+1}-1} \right\} + O\left(\frac{1}{(\log x)^A}\right), \\ \frac{1}{x} \sum_{n \leq x} \frac{\phi(n+3)\phi(n+2)\phi(n+1)}{(n+3)(n+2)(n+1)} &= w_2 \prod_{p > 2} \left\{ 1 - \frac{3}{p^2} \right\} + O\left(\frac{1}{(\log x)^A}\right), \end{aligned}$$

where w_2 is defined by (7) at $p = 2$ in which $g_j, j = 1, 2, 3$ are replaced by σ_a and ϕ respectively to the above sums.

Application 4. If ϕ, μ and $\sigma_a, a > 0$ are as above then as $x \rightarrow \infty$,

$$\begin{aligned} \frac{1}{x} \sum_{n \leq x} \frac{\phi(n+3)}{(n+3)} \frac{\phi(n+2)}{(n+2)} \mu(n+1) &= o(1), \\ \frac{1}{x} \sum_{n \leq x} \frac{\sigma_a(n+3)}{(n+3)^a} \frac{\sigma_a(n+2)}{(n+2)^a} \mu(n+1) &= o(1). \end{aligned}$$

Application 5. If ϕ, μ and $\sigma_a, a > 0$ are as above and under Assumption 1, we have as $x \rightarrow \infty$,

$$\begin{aligned} \frac{1}{x} \sum_{n \leq x} \frac{\phi(n+3)}{(n+3)} \mu(n+2) \mu(n+1) &= o(1), \\ \frac{1}{x} \sum_{n \leq x} \frac{\sigma_a(n+3)}{(n+3)^a} \mu(n+2) \mu(n+1) &= o(1). \end{aligned}$$

Application 6. Let $F_1(x) = x^2 + b, F_2(x) = x^2 + c, F_3(x) = x^2 + d, a > 0, 0 < t < 1$, where b, c, d are taken such that $F_j(x), j = 1, 2, 3$ satisfies the assumption of Theorem 2 and quadratic residue for all odd prime p . Then there exist a natural number γ such that for all $x \geq \gamma$,

$$\begin{aligned} \frac{1}{x} \sum_{n \leq x} \frac{\sigma_a(n^2 + b) \sigma_a(n^2 + c) \sigma_a(n^2 + d)}{(n^2 + b)^a (n^2 + c)^a (n^2 + d)^a} &= P'_1(\gamma) \prod_{p > \gamma} \left(1 + \frac{6}{p^{a+1} - 1} \right) + O\left(\frac{1}{(\log x)^t} \right) \\ \frac{1}{x} \sum_{n \leq x} \frac{\phi(n^2 + b) \phi(n^2 + c) \phi(n^2 + d)}{(n^2 + b)(n^2 + c)(n^2 + d)} &= P''_1(\gamma) \prod_{p > \gamma} \left(1 - \frac{6}{p^2} \right) + O\left(\frac{1}{(\log x)^t} \right), \end{aligned}$$

where $P'_1(\gamma)$ and $P''_1(\gamma)$ are defined by (8) in which $g_j, j = 1, 2, 3$ are replaced by σ_a and ϕ respectively.

Applications 7 and 8 are the behaviour of the distribution of the sum (2).

Application 7. Let f_1, f_2 and f_3 be real-valued additive functions and

$$(16) \quad \sum_{|f_j(p)| \leq 1} \frac{f_j^2(p)}{p} < \infty, j = 1, 2, 3,$$

$$(17) \quad \sum_{|f_j(p)| > 1} \frac{1}{p} < \infty, j = 1, 2, 3,$$

$$(18) \quad \sum_{j=1}^3 \sum_{|f_j(p)| \leq 1} \frac{f_j(p)}{p} < \infty,$$

Then the distribution functions

$$(19) \quad \frac{1}{[x]} \# \{n | n \leq x, f_1(n+3) + f_2(n+2) + f_3(n+1) \leq z\}$$

converge weakly towards a limit distribution ([3], Chapter III.2) as $x \rightarrow \infty$, and the characteristic function of this limit distribution is equal to

$$(20) \quad w_2 \prod_{p > 2} \left(1 - \frac{3}{p} + \left(1 - \frac{1}{p} \right) \sum_{m=1}^{\infty} \sum_{k=1}^3 \frac{\exp(it f_k(p^m))}{p^m} \right),$$

where w_2 is defined by (7) at $p = 2$ with g_k is replaced by $\exp(itf_k)$, $k = 1, 2, 3$.

Application 8. Let f_1, f_2 and f_3 be real-valued additive functions and $F_j(n)$, $j = 1, 2, 3$ are as above of degree $v_j \geq 2$. Assume that

$$(21) \quad \sum_{|f_j(p)| \leq 1} \frac{f_j^2(p)}{p} \varrho_j(p) < \infty, j = 1, 2, 3,$$

$$(22) \quad \sum_{|f_j(p)| > 1} \frac{\varrho_j(p)}{p} < \infty, j = 1, 2, 3,$$

$$(23) \quad \sum_{j=1}^3 \sum_{|f_j(p)| \leq 1} \frac{f_j(p)\varrho_j(p)}{p} < \infty,$$

$$(24) \quad f_j(p^m)\varrho_j(p^m) \rightarrow 0,$$

for $m = 1$, when $v_j \geq 2$ and for $m = 1, 2, \dots, v_j - 2$, when $v_j \geq 3$. Then the distribution functions

$$(25) \quad \frac{1}{[x]} \# \{n | n \leq x, f_1(F_1(n)) + f_2(F_2(n)) + f_3(F_3(n)) \leq z\}$$

converge weakly towards a limit distribution as $x \rightarrow \infty$, and the characteristic function of this limit distribution is equal to $P_1(\gamma)P_2(\gamma)$, where $P_1(\gamma)$ and $P_2(\gamma)$ are defined by (8) and (15) respectively with g_j is replaced by $\exp(itf_j)$, $j = 1, 2, 3$.

5. Proof of Theorem 2

We begin with some lemmas. The first lemma is required for the polynomial version of classical Turán-Kubilius inequality.

Lemma 1 ([8], Lemma 3). Let, $F(m)$ be arbitrary primitive polynomial of degree v with integer coefficients and with discriminant D . Let, $D \neq 0$. Then the number of solution of the congruence $F(m) \equiv 0 \pmod{p^\alpha}$ is $\varrho(p)$, when $p \nmid D$, and smaller than vD^2 when $p | D$.

Further, $\varrho(ab) = \varrho(a)\varrho(b)$, if $(a, b) = 1$.
and $\varrho(p^\alpha) \leq c$, c depends only on F .

Now we prove a polynomial version of classical Turán-Kubilius inequality.

Lemma 2. Let, $f(p^m)$ be the sequence of complex numbers for all $p \in \wp$, $m \geq 1$ and $F(n)$ is a polynomial as above of degree v . Then we have

$$\sum_{n \leq x} |K(F(n)) - A(x)| \ll xB(F(x)) + \sum_{m=1}^{v-1} \sum_{p^m \leq F(x)} |f(p^m)| \varrho(p^m) + \frac{x}{\log x},$$

where

$$K(n) := \sum_{p^m \parallel n} f(p^m), \quad A(x) := \sum_{p^m \leq x} \frac{f(p^m)\varrho(p^m)}{p^m},$$

$$B^2(x) := \sum_{p^m \leq x} \frac{|f(p^m)|^2 \varrho(p^m)}{p^m}.$$

Proof. We write $K(F(n)) = \sum_{p^m \parallel F(n)} f(p^m) = g(F(n)) + h(F(n))$,

where

$$g(y) = \sum_{\substack{p^m \parallel y \\ p^m \leq y}} f(p^m) \quad \text{and} \quad h(y) = \sum_{\substack{p^m \parallel y \\ p^m > y}} f(p^m).$$

Now,

$$\sum_{n \leq x} |K(F(n)) - A(x)| \leq \sum_{n \leq x} |g(F(n)) - A(x^{1/2})| + \sum_{n \leq x} |h(F(n))| + \sum_{n \leq x} |A(x^{1/2}) - A(x)|.$$

From Turán-Kubilius inequality ([1], Lemma 4.11), we have

$$\sum_{n \leq x} |g(F(n)) - A(x^{1/2})| \ll xB(x^{1/2}).$$

From Lemma 1 and Cauchy-Schwarz inequality, we have

$$\begin{aligned} |A(x) - A(x^{1/2})| &\leq \sum_{x^{1/2} < p^m \leq x} \frac{|f(p^m)| \varrho(p^m)}{p^m} \\ &\leq \left(\sum_{x^{1/2} < p^m \leq x} \frac{|f(p^m)|^2 \varrho(p^m)}{p^m} \right)^{1/2} \left(\sum_{x^{1/2} < p^m \leq x} \frac{\varrho(p^m)}{p^m} \right)^{1/2} = O(B(x)). \end{aligned}$$

Again by Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sum_{n \leq x} |h(F(n))| &= \sum_{n \leq x} \left| \sum_{\substack{p^m \parallel F(n) \\ p^m > x^{1/2}}} f(p^m) \right| \\ &\ll \sum_{x^{1/2} < p^m \leq F(x)} \frac{x|f(p^m)| \varrho(p^m)}{p^m} + \sum_{x^{1/2} < p^m \leq F(x)} |f(p^m)| \varrho(p^m) \\ &\ll x \left(\sum_{x^{1/2} < p^m \leq F(x)} \frac{|f(p^m)|^2 \varrho(p^m)}{p^m} \right)^{1/2} \left(\sum_{x^{1/2} < p^m \leq F(x)} \frac{\varrho(p^m)}{p^m} \right)^{1/2} \\ &+ \sum_{x^{1/2} < p^m \leq F(x)} |f(p^m)| \varrho(p^m) \ll xB(F(x)) + \sum_{m=1}^{v-1} \sum_{p^m \leq F(x)} |f(p^m)| \varrho(p^m) + \frac{x}{\log x}. \end{aligned}$$

Which proves the required Lemma. $\square$

The following lemma ensures the existence of γ in Theorem 2.

Lemma 3 ([9], Lemma 2.1). *If $F_1(m)$ and $F_2(m)$ are relatively prime polynomials with integer coefficients, then the congruence $F_1(m) \equiv 0 \pmod{a}$, $F_2(m) \equiv 0 \pmod{a}$ have common roots atmost for finitely many values of a .*

Define multiplicative functions g_{jr} and g_{jr}^* , $j = 1, 2, 3$ by

$$g_{jr}(p^m) = \begin{cases} g_j(p^m) & \text{if } p \leq r \\ 1 & \text{if } p > r. \end{cases} \quad \text{and} \quad g_{jr}^*(p^m) = \begin{cases} 1 & \text{if } p \leq r \\ g_j(p^m) & \text{if } p > r. \end{cases}$$

and multiplicative function h_{jr} , $j = 1, 2, 3$ by

$$h_{jr}(p^m) = \begin{cases} g_j(p^m) - g_j(p^{m-1}) & \text{if } p \leq r \\ 0 & \text{if } p > r. \end{cases}$$

so that, $g_{jr} = 1 * h_{jr}$, $j = 1, 2, 3$.

5.1. Proof of Theorem 2. We can write

$$\begin{aligned} M_x''(g_1, g_2, g_3) - P'(x) &= P'(r, x) \left(\frac{1}{x} \sum_{n \leq x} g_{1r}(F_1(n)) g_{2r}(F_2(n)) g_{3r}(F_3(n)) - P'(r) \right) \\ &+ \frac{1}{x} \sum_{n \leq x} g_{1r}(F_1(n)) g_{2r}(F_2(n)) g_{3r}(F_3(n)) \left(g_{1r}^*(F_1(n)) g_{2r}^*(F_2(n)) g_{3r}^*(F_3(n)) - P'(r, x) \right). \end{aligned}$$

So,

$$\begin{aligned} \left| M_x''(g_1, g_2, g_3) - P'(x) \right| &\leq \frac{1}{x} \left| \sum_{n \leq x} g_{1r}(F_1(n)) g_{2r}(F_2(n)) g_{3r}(F_3(n)) - P'(r) \right| \\ &+ \frac{1}{x} \sum_{n \leq x} \left| g_{1r}^*(F_1(n)) g_{2r}^*(F_2(n)) g_{3r}^*(F_3(n)) - P'(r, x) \right| := S_1 + S_2. \end{aligned}$$

Estimation of S_1 .

$$\begin{aligned} \frac{1}{x} \sum_{n \leq x} g_{1r}(F_1(n)) g_{2r}(F_2(n)) g_{3r}(F_3(n)) &= \frac{1}{x} \sum_{\substack{d_j \leq F_j(x) \\ j=1,2,3}} h_{1r}(d_1) h_{2r}(d_2) h_{3r}(d_3) \sum_{\substack{n \leq x \\ d_j | \bar{F}_j(x) \\ j=1,2,3}} 1 \\ &= \frac{1}{x} \sum_{d_1 \leq F_1(x)} \sum_{d_2 \leq F_2(x)} \sum_{d_3 \leq F_3(x)} h_{1r}(d_1) h_{2r}(d_2) h_{3r}(d_3) \frac{x}{[d_1, d_2, d_3]} \varrho(d_1, d_2, d_3) \\ &+ O\left(\frac{1}{x} \sum_{\substack{d_i \leq F_j(x) \\ j=1,2,3}} h_{1r}(d_1) h_{2r}(d_2) h_{3r}(d_3) \varrho(d_1, d_2, d_3) \right) := P'_1 + S_3. \end{aligned}$$

Now we observe that

$$\sum_{d_j=1}^{\infty} \frac{|h_{jr}(d_j)| \varrho_j(d_j)}{d_j} \leq \exp\left(c_{F_j} \sum_{p \leq r} \frac{1}{p} \right) \ll (\log r)^{c_{F_j}}$$

and for $0 < \alpha < 1$

$$\sum_{d_j=1}^{\infty} \frac{|h_{jr}(d_j)| \varrho_j(d_j)}{d_j^{1-\alpha}} \leq \prod_{p \leq r} \left(1 + \sum_{m=1}^{\infty} \frac{|h_{jr}(p^m)| \varrho_j(p^m)}{p^{m(1-\alpha)}} \right) \leq \exp\left(c_{F_j} \frac{r^\alpha}{\log r} \right).$$

We can say that,

$$\begin{aligned} S_3 &\ll \frac{1}{x} \sum_{\substack{d_j \leq F_j(x) \\ j=1,2,3}} |h_{1r}(d_1) h_{2r}(d_2) h_{3r}(d_3)| \varrho(d_1) \varrho(d_2) \varrho(d_3) \\ &\ll \frac{1}{x} (F_1(x) F_2(x) F_3(x))^{1-\alpha} \sum_{\substack{d_j=1 \\ j=1,2,3}}^{\infty} \frac{|h_{1r}(d_1)|}{d_1^{1-\alpha}} \frac{|h_{2r}(d_2)|}{d_2^{1-\alpha}} \frac{|h_{3r}(d_3)|}{d_3^{1-\alpha}} \varrho(d_1) \varrho(d_2) \varrho(d_3) \\ &\ll \frac{1}{x} (F_1(x) F_2(x) F_3(x))^{1-\alpha} \exp(c_F \frac{r^\alpha}{\log r}). \end{aligned}$$

Now,

$$\begin{aligned} P'_1 &= P'(r) + O\left(\sum_{k=1}^3 \sum_{\substack{d_j=1 \\ j=1,2,3 \\ d_k > F_k(x)}}^{\infty} \frac{|h_{1r}(d_1)h_{2r}(d_2)h_{3r}(d_3)|}{[d_1, d_2, d_3]} \varrho(d_1, d_2, d_3)\right) \\ &:= P'(r) + S_4. \end{aligned}$$

Again from the above observations, we have

$$\begin{aligned} S_4 &\ll \sum_{k=1}^3 \sum_{\substack{d_j=1 \\ j=1,2,3 \\ d_k > F_k(x)}}^{\infty} \frac{|h_{1r}(d_1)h_{2r}(d_2)h_{3r}(d_3)|}{d_1 d_2 d_3} \varrho(d_1) \varrho(d_2) \varrho(d_3) \\ &\ll (F_1(x)^{-\alpha} + F_2(x)^{-\alpha} + F_3(x)^{-\alpha}) \exp\left(c_F \frac{r^\alpha}{\log r}\right). \end{aligned}$$

Estimation of S_2 . Let

$$N'_r = \left\{ n \leq x \mid \exists k \in \{1, 2, 3\} \text{ and } \exists p > r \text{ such that } p^m \parallel F_k(n), |1 - g_k(p^m)| > \frac{1}{2} \right\}.$$

Decompose S_2 into two sums

$$\begin{aligned} S_2 &= \frac{1}{x} \sum_{n \in N'_r} \left| g_{1r}^*(F_1(n)) g_{2r}^*(F_2(n)) g_{3r}^*(F_3(n)) - P'(r, x) \right| \\ &\quad + \frac{1}{x} \sum_{n \notin N'_r} \left| g_{1r}^*(F_1(n)) g_{2r}^*(F_2(n)) g_{3r}^*(F_3(n)) - P'(r, x) \right| \\ &:= S_5 + S_6. \end{aligned}$$

Let us put,

$$\eta_j(p) := \sum_{m=1}^{\infty} \frac{(g_j(p^m) - g_j(p^{m-1})) \varrho_j(p^m)}{p^m}, j = 1, 2, 3.$$

From Lemma 1, we have

$$|\eta_j(p)| \leq 2c_{F_j} \frac{1}{p-1} \leq \frac{1}{6} \text{ if } p \geq 1 + 12c_{F_j} := p_j.$$

Let, $p_4 := \max(p_1, p_2, p_3)$. So, if $r \geq p_4$, then

$$\begin{aligned} P'(r, x) &= \prod_{r < p \leq x} (1 + \eta_1(p) + \eta_2(p) + \eta_3(p)) \\ &= \exp\left(\sum_{r < p \leq x} (\eta_1(p) + \eta_2(p) + \eta_3(p) + O(|\eta_1(p)|^2 + |\eta_2(p)|^2 + |\eta_3(p)|^2))\right) \\ &= \exp\left(\sum_{r < p \leq x} \sum_{j=1}^3 \frac{(g_j(p) - 1) \varrho_j(p)}{p} + O\left(\sum_{r < p \leq x} \frac{1}{p^2}\right)\right) \ll 1. \end{aligned}$$

Without loss of generality we may assume that, $r \geq p_4$.

Now

$$\begin{aligned}
S_5 &\ll \frac{1}{x} \sum_{n \in N'_r} 1 \ll \frac{1}{x} \sum_{\substack{p^m \leq F_j(x) \\ |1-g_j(p^m)| > 1/2 \\ p > r}} \left(\frac{x \varrho_j(p^m)}{p^m} + \varrho_j(p^m) \right) \\
&\ll \frac{1}{x} \sum_{\substack{p^m \leq F_j(x) \\ |1-g_j(p^m)| > 1/2 \\ p > r}} \frac{x \varrho_j(p^m)}{p^m} + \frac{1}{x} \sum_{\substack{p^m \leq F_j(x) \\ |1-g_j(p^m)| > 1/2 \\ p > r}} \varrho_j(p^m) \\
&\ll \sum_{r < p \leq F_j(x)} \frac{|1-g_j(p)| \varrho_j(p)}{p} + \sum_{p > r} \frac{1}{p^2} + \frac{1}{x} \sum_{\substack{p^m \leq F_j(x) \\ p > r, m < v_j}} |1-g_j(p^m)| \varrho_j(p^m) + \frac{1}{\log x} \\
&\ll S'(r, x) + T(x) + (r \log r)^{-1} + \frac{1}{x} C(r) + \frac{1}{\log x}.
\end{aligned}$$

Since we know that if $\operatorname{Re}(u) \leq 0, \operatorname{Re}(v) \leq 0$, then

$$(26) \quad |\exp(u) - \exp(v)| \leq |u - v| \quad \text{and} \quad \text{for } |z| \leq 1, |\arg(z)| \leq \frac{\pi}{2}$$

$$(27) \quad \log(1+z) = z + O(|z|^2).$$

we have,

$$\begin{aligned}
S_6 &\ll \frac{1}{x} \sum_{n \leq x} \sum_{j=1}^3 \left| \sum_{\substack{p^m \parallel F_j(n) \\ p > r}} (g_j(p^m) - 1) - \sum_{\substack{p^m \leq x \\ p > r}} \frac{g_j(p^m) - 1}{p^m} \varrho_j(p^m) \right| \\
&\quad + \frac{1}{x} \sum_{n \leq x} \left| \sum_{\substack{p^m \leq x \\ p > r}} \sum_{j=1}^3 \frac{(g_j(p^m) - 1) \varrho_j(p^m)}{p^m} - \log P'(r, x) \right| \\
&\quad + O\left(\frac{1}{x} \sum_{n \leq x} \sum_{j=1}^3 \sum_{\substack{p^m \parallel F_j(x) \\ p > r}} |g_j(p^m) - 1|^2 \right) := S_{61} + S_{62} + S_{63}.
\end{aligned}$$

From Lemma 2, we have

$$\begin{aligned}
S_{61} &\ll \sum_{j=1}^3 \left(\sum_{\substack{p^m \leq F_j(x) \\ p > r}} \frac{|g_j(p^m) - 1|^2 \varrho_j(p^m)}{p^m} \right)^{1/2} + \frac{1}{x} C(r) + \frac{1}{\log x} \\
&\ll (S'(r, x))^{1/2} + (T(x))^{1/2} + (r \log r)^{-1/2} + \frac{1}{x} C(r) + \frac{1}{\log x}.
\end{aligned}$$

$$\begin{aligned}
S_{62} &= \left| \sum_{j=1}^3 \sum_{r < p \leq x} \frac{(g_j(p) - 1) \varrho_j(p)}{p} + O\left(\sum_{r < p} \frac{1}{p^2} \right) - \sum_{r < p \leq x} \log w'_p \right| \\
&= \left| \sum_{j=1}^3 \sum_{r < p \leq x} \frac{(g_j(p) - 1) \varrho_j(p)}{p} + O\left(\sum_{r < p} \frac{1}{p^2} \right) - \sum_{r < p \leq x} \sum_{j=1}^3 \frac{(g_j(p) - 1) \varrho_j(p)}{p} \right| \\
&\ll \sum_{p > r} \frac{1}{p^2} \ll (r \log r)^{-1}.
\end{aligned}$$

$$\begin{aligned}
S_{63} &\ll \sum_{j=1}^3 \sum_{\substack{p^m \leq F_j(x) \\ p > r}} \frac{|g_j(p^m) - 1|^2 \varrho_j(p^m)}{p^m} + \frac{1}{x} C(r) + \frac{1}{\log x} \\
&\ll S'(r, x) + T(x) + (r \log r)^{-1} + \frac{1}{x} C(r) + \frac{1}{\log x}.
\end{aligned}$$

Combining all these estimates for all $1 - \frac{1}{v_1 + v_2 + v_3} < \alpha < 1$, we have

$$\begin{aligned}
M_x''(g_1, g_2, g_3) - P'(x) &\ll (F_1(x)F_2(x)F_3(x))^{1-\alpha} \exp\left(c \frac{r^\alpha}{\log r}\right) + (S'(r, x))^{1/2} \\
&\quad + (T(x))^{1/2} + (r \log r)^{-1/2} + \frac{1}{x} C(r) + \frac{1}{\log x}.
\end{aligned}$$

which proves the required Theorem.

6. Proof of Theorem 3 and 4

We begin with some lemmas. The first lemma will be used to prove Theorem 3.

Lemma 4 ([5], Theorem 1). *For any given $K > 0$,*

$$\sum_{n \leq x} \mu(n) \exp(2\pi i n \theta) = O\left(\frac{x}{(\log x)^K}\right),$$

uniformly in θ .

The next lemmas will be used to prove Theorem 4.

Lemma 5 ([2], Theorem 2). *Let g be a multiplicative function whose modulus does not exceed 1. Then there is a real τ , $|\tau| \leq (\log x)^{1/19}$, such that*

$$(28) \quad \sum_{\substack{n \leq x \\ (n, D)=1}} g(n) = \theta_\tau(D) \sum_{n \leq x} g(n) + O\left(\frac{x(\log \log 3d)^2}{(\log x)^{1/19}}\right)$$

holds uniformly for $x \geq 2$ and odd integers D . If, in addition, the condition (11) is satisfied then (31) holds for even integers as well. For real-valued g we may set $\tau = 0$.

The following lemma is a special case of a theorem of Wolke [7].

Lemma 6 ([7], Theorem 1). *Let g be as above and g is a good function. Then for given any $A > 0$ there is a corresponding $A_1 > 0$, possibly depending on g , such that for $x \geq 2$ and $Q = x^{1/2}(\log x)^{-A_1}$, we have*

$$\sum_{d \leq Q} \max_{(l, d)=1} \max_{u \leq x} \left| \sum_{\substack{n \leq u \\ n \equiv l \pmod{d}}} g(n) - \frac{1}{\phi(d)} \sum_{\substack{n \leq u \\ (n, d)=1}} g(n) \right| \ll \frac{x}{(\log x)^A}$$

In case $-\tau \in \mathbb{N}$ or $\tau = 0$ then

$$\sum_{d \leq Q} \max_l \max_{u \leq x} \left| \sum_{\substack{n \leq u \\ n \equiv l \pmod{d}}} g(n) \right| \ll \frac{x}{(\log x)^A}.$$

The following lemma is a two dimensional version of standard Cauchy-Schwarz inequality:

Lemma 7. *If x_j, x_k and c_{jk} are non-negative real numbers, then*

$$\sum_{j \leq y} \sum_{k \leq y} x_j x_k c_{jk} \leq \left(\sum_{j \leq y} \sum_{k \leq y} x_j^2 x_k^2 c_{jk} \right)^{1/2} \left(\sum_{j \leq y} \sum_{k \leq y} c_{jk} \right)^{1/2}.$$

Proof. By applying standard Cauchy-Schwarz over k , we have

$$\sum_{j \leq y} \sum_{k \leq y} x_j x_k c_{jk} \leq \sum_{j \leq y} \left(\sum_{k \leq y} x_j^2 x_k^2 c_{jk} \right)^{1/2} \left(\sum_{k \leq y} c_{jk} \right)^{1/2} := \sum_{j \leq y} a_j b_j.$$

Again by applying cauchy-Schwarz over j , we have

$$\sum_{j \leq y} a_j b_j \leq \left(\sum_{j \leq y} a_j^2 \right)^{1/2} \left(\sum_{j \leq y} b_j^2 \right)^{1/2} = \left(\sum_{j \leq y} \sum_{k \leq y} x_j^2 x_k^2 c_{jk} \right)^{1/2} \left(\sum_{j \leq y} \sum_{k \leq y} c_{jk} \right)^{1/2}.$$

□

6.1. Proof of Theorem 3. Let us put,

$$R(r, x) = \prod_{r < p \leq x} \left\{ 1 - \frac{2}{p} + \left(1 - \frac{1}{p} \right) \sum_{m=1}^{\infty} \frac{g_1(p^m) + g_2(p^m)}{p^m} \right\}.$$

It is easy to see that, $|R(r, x)| \leq 1$. Therefore,

$$\begin{aligned} M'_x(g_1, g_2, \mu) &= R(r, x) \frac{1}{x} \sum_{n \leq x} g_{1r}(n+3) g_{2r}(n+2) \mu(n+1) \\ &+ \frac{1}{x} \sum_{n \leq x} g_{1r}(n+3) g_{2r}(n+2) \mu(n+1) (g_{1r}^*(n+3) g_{2r}^*(n+2) - R(r, x)). \end{aligned}$$

So,

$$\begin{aligned} x \left| M'_x(g_1, g_2, \mu) \right| &\leq \left| \sum_{n \leq x} g_{1r}(n+3) g_{2r}(n+2) \mu(n+1) \right| + \sum_{n \leq x} \left| g_{1r}^*(n+3) g_{2r}^*(n+2) - R(r, x) \right| \\ &:= T_1 + T_2. \end{aligned}$$

Estimation of T_1 .

$$\begin{aligned} \sum_{n \leq x} g_{1r}(n+3) g_{2r}(n+2) \mu(n+1) &= \sum_{n \leq x} \sum_{d_1 | n+3} \sum_{d_2 | n+2} h_{1r}(d_1) h_{2r}(d_2) \mu(n+1) \\ &= \sum_{d_1 \leq x+3} \sum_{d_2 \leq x+2} h_{1r}(d_1) h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ d_1 | n+2 \\ d_2 | n+1}} \mu(n) = \sum_{d_1 \leq x+3} \sum_{\substack{d_2 \leq x+2 \\ (d_1, d_2)=1}} h_{1r}(d_1) h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ n \equiv v(d_1 d_2)}} \mu(n) \\ &= \sum_{d_1 \leq y} h_{1r}(d_1) \sum_{\substack{d_2 \leq y \\ (d_1, d_2)=1}} h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ n \equiv v(d_1 d_2)}} \mu(n) + \sum_{k=1}^2 \sum_{\substack{d_j \leq x+4-j \\ j=1,2 \\ d_k > y \\ (d_1, d_2)=1}} h_{1r}(d_1) h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ n \equiv v(d_1 d_2)}} \mu(n), \end{aligned}$$

where v is unique solution of the system of linear congruence $n \equiv -2(d_1), n \equiv -1(d_2), 0 \leq v \leq d_1 d_2 - 1$ and $y := \log x$.

So,

$$T_1 \ll \sum_{\substack{d_j \leq y \\ j=1,2}} |h_{1r}(d_1)h_{2r}(d_2)| \left| \sum_{\substack{n \leq x+1 \\ n \equiv v(d_1 d_2)}} \mu(n) \right| + \sum_{k=1}^2 \sum_{\substack{d_j \leq x+4-j \\ j=1,2 \\ d_k > y}} |h_{1r}(d_1)h_{2r}(d_2)| \left(\frac{x}{d_1 d_2} + 1 \right) \\ := T_{11} + T_{12}.$$

From Lemma 4 we have,

$$\sum_{\substack{n \leq x+1 \\ n \equiv v(d_1 d_2)}} \mu(n) = \sum_{n \leq x+1} \mu(n) \frac{1}{d_1 d_2} \sum_{l=1}^{d_1 d_2} \exp\left(\frac{(n-v)l}{d_1 d_2}\right) \\ = \frac{1}{d_1 d_2} \sum_{l=1}^{d_1 d_2} \exp\left(\frac{-vl}{d_1 d_2}\right) \sum_{n \leq x+1} \mu(n) \exp\left(\frac{nl}{d_1 d_2}\right) \ll \frac{x}{(\log x)^K}.$$

So,

$$T_{11} \ll \frac{x}{(\log x)^K} \sum_{d_1 \leq y} \sum_{d_2 \leq y} |h_{1r}(d_1)h_{2r}(d_2)| \ll \frac{x}{(\log x)^K} y^4 \sum_{d_1=1}^{\infty} \sum_{d_2=1}^{\infty} \frac{|h_{1r}(d_1)h_{2r}(d_2)|}{d_1^2 d_2^2} \\ \ll \frac{x}{\log x}, \text{ if } K \geq 5.$$

Now from the following two estimations

$$(29) \quad \sum_{d_j=1}^{\infty} \frac{|h_{jr}(d_j)|}{d_j^\alpha} = \prod_{p \leq r} \left(1 + \sum_{m=1}^{\infty} \frac{|h_{jr}(p^m)|}{p^{m\alpha}} \right) \leq \prod_{p \leq r} \left(1 + \frac{2}{p^\alpha - 1} \right) \\ \leq \exp\left(c_1 \sum_{p \leq r} \frac{1}{p^\alpha}\right) \leq \exp\left(c_2 \frac{r^{1-\alpha}}{\log r}\right).$$

and

$$(30) \quad \sum_{d_j=1}^{\infty} \frac{|h_{jr}(d_j)|}{d_j} \leq \exp\left(c_3 \sum_{p \leq r} \frac{1}{p}\right) \ll (\log r)^{c_4}$$

We have,

$$T_{12} \ll x \sum_{\substack{d_j \leq x+4-j \\ j=1,2 \\ d_k > y}} \frac{|h_{1r}(d_1)h_{2r}(d_2)|}{d_1 d_2} + \sum_{\substack{d_i \leq x+4-j \\ j=1,2 \\ d_k > y}} |h_{1r}(d_1)h_{2r}(d_2)| \\ \ll xy^{-\gamma} \exp\left(c_2 \frac{r^\gamma}{\log r}\right) (\log r)^{c_4} + x^{2\alpha} \exp\left(2c_2 \frac{r^{1-\alpha}}{\log r}\right) \\ \ll xy^{-\gamma} \exp\left(c_5 \frac{r^\gamma}{\log r}\right) + x^{2\alpha} \exp\left(2c_2 \frac{r^{1-\alpha}}{\log r}\right).$$

Taking $1 - \alpha = \gamma = \frac{2}{3}$ we have,

$$T_{12} \ll xy^{-\frac{2}{3}} \exp\left(c_5 \frac{r^{2/3}}{\log r}\right) + x^{2/3} \exp\left(2c_2 \frac{r^{2/3}}{\log r}\right).$$

Putting $r = (\log \log x)^{3/2}$ we have

$$T_{12} \ll \frac{x}{y^{2/3}} (\log x)^{1/6} + xx^{-1/3} (\log x)^{1/6} \ll \frac{x}{(\log x)^{1/2}}.$$

So as $x \rightarrow \infty$,

$$T_1 = o(x).$$

Estimation of T_2 . To get an estimate of T_2 we will use an technique of R. Warlimont [6].

Now, let

$$N_r = \left\{ n \leq x \mid \exists j \in \{1, 2\} \text{ and } \exists p > r \text{ such that } p^m \parallel n + 4 - j, |1 - g_j(p^m)| > \frac{1}{2} \right\}$$

Decompose T_2 into two sums

$$\begin{aligned} T_2 &= \frac{1}{x} \sum_{n \in N_r} |g_{1r}^*(n+3)g_{2r}^*(n+2) - R(r, x)| + \frac{1}{x} \sum_{n \notin N_r} |g_{1r}^*(n+3)g_{2r}^*(n+2) - R(r, x)| \\ &:= T_5 + T_6. \end{aligned}$$

Now,

$$T_5 \ll \frac{1}{x} \sum_{j=1}^2 \sum_{r < p \leq x+4-j} \frac{x+4-j}{p} |1 - g_j(p)|^2 + \sum_{j=1}^2 \sum_{p > r} \sum_{m \geq 2} \frac{1}{p^m} \ll S''(r, x) + (r \log r)^{-1}$$

From (26) and (27), we have

$$\begin{aligned} T_6 &\leq \frac{1}{x} \sum_{j=1}^2 \sum_{n \leq x} \left| \sum_{\substack{p^m \parallel n+4-j \\ p > r}} (g_j(p^m) - 1) - \sum_{\substack{p^m \leq x \\ p > r}} \frac{g_j(p^m) - 1}{p^m} \right| + \frac{1}{x} \left| \sum_{\substack{p^m \leq x \\ p > r}} \sum_{j=1}^2 \frac{g_j(p^m) - 1}{p^m} - \log R(r, x) \right| \\ &+ O\left(\frac{1}{x} \sum_{n \leq x} \sum_{j=1}^2 \sum_{\substack{p^m \parallel n+4-j \\ p > r}} |g_j(p^m) - 1|^2\right) := T_7 + T_8 + T_9. \end{aligned}$$

Now by Cauchy-Schwarz inequality and Turán-Kubilius inequality ([1], Lemma 4.4), we have

$$\begin{aligned} T_7 &\ll \left(\sum_{j=1}^2 \sum_{\substack{p^m \leq x+4-j \\ p > r}} |g_j(p^m) - 1|^2 \right)^{\frac{1}{2}} + \frac{1}{x} \ll \left(\sum_{j=1}^2 \sum_{r < p \leq x+4-j} |g_j(p) - 1|^2 \right)^{\frac{1}{2}} + \left(\sum_{p > r} \frac{1}{p^2} \right)^{\frac{1}{2}} + \frac{1}{x} \\ &\ll \left(S''(r, x) \right)^{\frac{1}{2}} + (r \log r)^{-\frac{1}{2}} + x^{-1}. \end{aligned}$$

Now similar to estimation of S_{62} , we have

$$T_8 \ll \left| \sum_{r < p \leq x} \sum_{j=1}^2 \frac{g_j(p^m) - 1}{p^m} - \log R(r, x) \right| \ll \sum_{p > r} \frac{1}{p^2} \ll (r \log r)^{-1}.$$

and

$$T_9 \ll \frac{1}{x} \left\{ \sum_{j=1}^2 \sum_{\substack{p^m \leq x+4-j \\ p > r}} \frac{|g_j(p^m) - 1|^2}{p^m} \right\} \ll S''(r, x) + (r \log r)^{-1}.$$

Combining above calculations, we have

$$T_2 \ll (r \log r)^{-1/2} + (S(r, x))^{1/2} + x^{-1}.$$

By the above choice of r and from (10) we have as $x \rightarrow \infty$,

$$T_2 = o(x).$$

Which proves the required Theorem.

6.2. Proof of Theorem 4.

$$\begin{aligned} R &:= M'_x(g_1, g_2, g_3) - M_x(g_3)P''(r, x)Q(r) = P''(r, x) \left(M'_x(g_{1r}, g_{2r}, g_3) - M_x(g_3)Q(r) \right) \\ &+ \frac{1}{x} \sum_{n \leq x} g_{1r}(n+3)g_{2r}(n+2)g_3(n+1) \left(g_{1r}^*(n+3)g_{2r}^*(n+2) - P''(r, x) \right) \end{aligned}$$

It is easy to see that $|P''(r, x)| \leq 1$. Therefore

$$\begin{aligned} R &\ll \left| M'_x(g_{1r}, g_{2r}, g_3) - M_x(g_3)Q(r) \right| + \frac{1}{x} \sum_{n \leq x} \left| g_{1r}^*(n+3)g_{2r}^*(n+2) - P''(r, x) \right| \\ &:= U_1 + U_2. \end{aligned}$$

Estimation of U_1 .

$$\begin{aligned} M'_x(g_{1r}, g_{2r}, g_3) &= \frac{1}{x} \sum_{n \leq x} \sum_{d_1 | n+3} \sum_{d_2 | n+2} h_{1r}(d_1)h_{2r}(d_2)g_3(n+1) \\ &= \frac{1}{x} \sum_{d_1 \leq x+3} h_{1r}(d_1) \sum_{\substack{d_2 \leq x+2 \\ (d_1, d_2)=1}} h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ n \equiv v(d_1 d_2)}} g_3(n) \\ &= \frac{1}{x} \sum_{\substack{d_j \leq y \\ j=1,2 \\ (d_1, d_2)=1}} h_{1r}(d_1)h_{2r}(d_2) \sum_{\substack{2 \leq n \leq x+1 \\ n \equiv v(d_1 d_2)}} g_3(n) + \frac{1}{x} \sum_{k=1}^2 \sum_{\substack{d_j \leq x+4-j \\ j=1,2 \\ d_k > y}} h_{1r}(d_1)h_{2r}(d_2) \left(\frac{x}{d_1 d_2} + 1 \right) \\ &:= P_2 + U_{11}, \end{aligned}$$

where v is the unique solution of the system $n \equiv -2(d_1), n \equiv -1(d_2)$ and $y := x^{1/4}(\log x)^{-\frac{\beta}{2}}$.

From (29), (30), for $0 < \alpha, \gamma < 1$, we have

$$\begin{aligned} U_{11} &\ll y^{-\gamma} \exp\left(c_6 \frac{r^\gamma}{\log r}\right) + x^{1-2\alpha} \exp\left(2c_2 \frac{r^\alpha}{\log r}\right) \\ &\ll x^{-\gamma/4}(\log x)^{\frac{\gamma\beta}{2}} \exp\left(c_6 \frac{r^\gamma}{\log r}\right) + x^{1-2\alpha} \exp\left(2c_2 \frac{r^\alpha}{\log r}\right). \end{aligned}$$

$$\begin{aligned} P_2 &= \frac{1}{x} \sum_{d_1 \leq y} \sum_{\substack{d_2 \leq y \\ (d_1, d_2)=1}} \frac{h_{1r}(d_1)h_{2r}(d_2)}{\phi(d_1 d_2)} \sum_{\substack{n \leq x \\ (n, d_1 d_2)=1}} g_3(n) + O\left(\frac{1}{x} \sum_{\substack{d_j \leq y \\ j=1,2}} |h_{1r}(d_2)h_{2r}(d_2)|\right) \\ &+ O\left(\frac{1}{x} \sum_{\substack{d_j \leq y \\ j=1,2}} |h_{1r}(d_1)h_{2r}(d_2)| \left| \sum_{\substack{n \leq x \\ n \equiv v(d_1 d_2)}} g_3(n) - \frac{1}{\phi(d_1 d_2)} \sum_{\substack{n \leq x \\ (n, d_1 d_2)=1}} g_3(n) \right| \right) := P_3 + U_{12} + U_{13}. \end{aligned}$$

By Lemma 6 and Lemma 7, we have

$$\begin{aligned}
U_{13} &\ll \frac{1}{x} \left(\sum_{l \leq y^2} \left| \sum_{\substack{n \leq x \\ n \equiv v(l)}} g_3(n) - \frac{1}{\phi(l)} \sum_{\substack{n \leq x \\ (n, l) = 1}} g_3(n) \right| \right)^{1/2} \\
&\quad \left(\sum_{\substack{d_j \leq y \\ j=1,2}} |h_{1r}(d_1)|^2 |h_{2r}(d_2)|^2 \left| \sum_{\substack{n \leq x \\ n \equiv v(d_1 d_2)}} g_3(n) - \frac{1}{\phi(d_1 d_2)} \sum_{\substack{n \leq x \\ (n, d_1 d_2) = 1}} g_3(n) \right| \right)^{1/2} \\
&\ll \frac{1}{(\log x)^{A/2}} \left(\sum_{\substack{d_j \leq y \\ j=1,2}} \frac{|h_{1r}(d_1)|^2 |h_{2r}(d_2)|^2}{\phi(d_1) \phi(d_2)} \right)^{1/2}.
\end{aligned}$$

Since

$$\sum_{d \leq y} \frac{|h(d)|^2}{\phi(d)} \leq \exp \left(c_7 \sum_{p \leq r} \frac{1}{p} \right) \leq (\log r)^{c_8},$$

we have

$$U_{13} \ll \frac{(\log r)^{c_8}}{(\log x)^{A/2}}.$$

Now from (29) we get

$$U_{12} \ll \frac{y^{2(1-\alpha)}}{x} \sum_{\substack{d_j=1 \\ j=1,2}}^{\infty} \frac{|h_{1r}(d_1) h_{2r}(d_2)|}{d_1^{1-\alpha} d_2^{1-\alpha}} \ll x^{-\frac{1}{2}(1+\alpha)} (\log x)^{-\beta(1-\alpha)} \exp \left(2c_2 \frac{r^\alpha}{\log r} \right).$$

Using Lemma 5, we get

$$\begin{aligned}
P_3 &= \frac{1}{x} \sum_{d_1 \leq y} \sum_{\substack{d_2 \leq y \\ (d_1, d_2) = 1}} \frac{h_{1r}(d_1) h_{2r}(d_2)}{\phi(d_1 d_2)} \theta_\tau(d_1 d_2) \sum_{n \leq x} g_3(n) \\
&\quad + O \left(\sum_{\substack{d_j \leq y \\ j=1,2 \\ (d_1, d_2) = 1}} \frac{|h_{1r}(d_1) h_{2r}(d_2)|}{\phi(d_1) \phi(d_2)} \frac{(\log \log 3d_1 d_2)^2}{(\log x)^{1/19}} \right) := P_4 + U_{14}.
\end{aligned}$$

Now,

$$\begin{aligned}
&\sum_{d_1 \leq y} \sum_{\substack{d_2 \leq y \\ (d_1, d_2) = 1}} \frac{h_{1r}(d_1) h_{2r}(d_2)}{\phi(d_1) \phi(d_2)} \theta_\tau(d_1) \theta_\tau(d_2) = \sum_{d_1=1}^{\infty} \sum_{\substack{d_2=1 \\ (d_1, d_2) = 1}}^{\infty} \frac{h_{1r}(d_1) h_{2r}(d_2)}{\phi(d_1) \phi(d_2)} \theta_\tau(d_1) \theta_\tau(d_2) \\
&\quad + O \left(\sum_{k=1}^2 \sum_{\substack{d_j \leq y \\ j=1,2 \\ d_k > y}} \frac{h_{1r}(d_1) h_{2r}(d_2)}{\phi(d_1) \phi(d_2)} \theta_\tau(d_1) \theta_\tau(d_2) \right) := P_5 + U_{15}.
\end{aligned}$$

Now from the following two estimation

$$\sum_{d > y} \frac{|h_{jr}(d) \theta_\tau(d)|}{\phi(d)} \leq y^{-\alpha} \exp \left(c_9 \sum_{p \leq r} \frac{1}{p^{1-\alpha}} \right) \ll x^{-\frac{1-\alpha}{4}} (\log x)^{\frac{\beta(1-\alpha)}{2}} \exp \left(c_{10} \frac{r^\alpha}{\log r} \right).$$

and

$$\sum_{d=1}^{\infty} \frac{|h_{jr}(d)\theta_{\tau}(d)|}{\phi(d)} \leq \exp\left(c_{11} \sum_{p \leq r} \frac{1}{p}\right) \leq (\log r)^{c_{12}}.$$

we can sat that

$$U_{15} \ll x^{-\frac{1-\alpha}{4}} (\log x)^{\frac{\beta(1-\alpha)}{2}} \exp\left(c_{13} \frac{r^{\alpha}}{\log r}\right).$$

$$\begin{aligned} P_5 &= \prod_{p \leq r} \left(1 + \sum_{m=1}^{\infty} \frac{(h_1(p^m) + h_2(p^m)) \theta_{\tau}(p^m)}{\phi(p^m)}\right) = \prod_{p \leq r} \left(1 - \frac{2\theta_{\tau}(p)}{p-1} + \theta_{\tau}(p) \sum_{m=1}^{\infty} \frac{g_1(p^m) + g_2(p^m)}{p^m}\right) \\ &= Q(r). \end{aligned}$$

Since

$$\begin{aligned} \sum_{d=1}^{\infty} \frac{|h_{jr}(d)| \log \log d}{\phi(d)} &= \prod_{p \leq r} \left(1 + \sum_{\alpha=1}^{\infty} \frac{|h_{jr}(p^{\alpha})| \alpha \log \log p}{\phi(p^{\alpha})}\right) \ll \prod_{p \leq r} \left(1 + \frac{2p \log \log p}{(p-1)^2}\right) \\ &\ll \exp\left(c_{14} \sum_{p \leq r} \frac{\log \log p}{p}\right) \ll \exp(c_{15} (\log \log r)^2), \end{aligned}$$

then we can say that

$$U_{14} \ll \frac{\exp(2c_{15} (\log \log r)^2)}{(\log x)^{1/19}}.$$

From the similar calculation as Estimation of T_2 , we have

$$U_2 \ll (S''(r, x))^{1/2} + (r \log r)^{-1/2}.$$

Combining these results, we get

$$\begin{aligned} R &\ll \left(x^{-\frac{\gamma}{4}} (\log x)^{\gamma\beta} + x^{\frac{\alpha-1}{4}} (\log x)^{\frac{\beta(1-\alpha)}{2}}\right) \exp\left(c_{16} \frac{r^{\gamma}}{\log r}\right) + x^{1-2\alpha} \exp\left(c_{17} \frac{r^{\alpha}}{\log r}\right) + \frac{(\log r)^{c_8}}{(\log x)^{\frac{4}{2}}} \\ &+ x^{-\frac{(1+\alpha)}{2}} (\log x)^{\beta(1-\alpha)} \exp\left(c_{18} \frac{r^{\alpha}}{\log r}\right) + \frac{\exp(2c_{15} (\log \log r)^2)}{(\log x)^{1/19}} + (S''(r, x))^{\frac{1}{2}} + (r \log r)^{-\frac{1}{2}}. \end{aligned}$$

By choosing $\alpha = \gamma$, we get the required theorem.

7. Proof of Theorem 5

Let us put,

$$T(r, x) = \prod_{r < p \leq x} \left\{1 - \frac{1}{p} + \left(1 - \frac{1}{p}\right) \sum_{m=1}^{\infty} \frac{g_1(p^m)}{p^m}\right\}.$$

Now,

$$\begin{aligned} \sum_{n \leq x} g_1(n+3) \mu(n+2) \mu(n+1) &= T(r, x) \sum_{n \leq x} g_{1r}(n+3) \mu(n+2) \mu(n+1) \\ &+ \sum_{n \leq x} g_{1r}(n+3) \mu(n+2) \mu(n+1) (g_{1r}^*(n+3) - T(r, x)). \end{aligned}$$

It is easy to see that $|T(r, x)| \leq 1$. Therefore,

$$\begin{aligned} \left|\sum_{n \leq x} g_1(n+3) \mu(n+2) \mu(n+1)\right| &\leq \left|\sum_{n \leq x} g_{1r}(n+3) \mu(n+2) \mu(n+1)\right| + \sum_{n \leq x} |g_{1r}^*(n+3) - T(r, x)| \\ &:= V_1 + V_2. \end{aligned}$$

Estimation of V_1 .

$$\begin{aligned} V_1 &= \left| \sum_{n \leq x} \sum_{d|n+3} h_{1r}(d) \mu(n+2) \mu(n+1) \right| = \left| \sum_{d \leq x+3} h_{1r}(d) \sum_{\substack{n \leq x \\ d|n+3}} \mu(n+2) \mu(n+1) \right| \\ &\leq \left| \sum_{d \leq y} h_{1r}(d) \sum_{\substack{n \leq x \\ n \equiv -3(d)}} \mu(n+2) \mu(n+1) \right| + \left| \sum_{y < d \leq x+3} h_{1r}(d) \sum_{\substack{n \leq x \\ n \equiv -3(d)}} \mu(n+2) \mu(n+1) \right| := V_{11} + V_{12}, \end{aligned}$$

where $y := \log x$.

Under Assumption 1, we have

$$\begin{aligned} \sum_{\substack{n \leq x \\ n \equiv -3(d)}} \mu(n+2) \mu(n+1) &= \sum_{n \leq x} \mu(n+2) \mu(n+1) \frac{1}{d} \sum_{l=1}^d \exp\left(\frac{(n-3)l}{d}\right) \\ &= \frac{1}{d} \sum_{l=1}^d \exp\left(\frac{-3l}{d}\right) \sum_{n \leq x} \mu(n+2) \mu(n+1) \exp\left(\frac{nl}{d}\right) \ll \frac{x}{(\log x)^A}. \end{aligned}$$

By choosing $A = 3$ we have,

$$V_{11} \ll \frac{x}{(\log x)^3} \sum_{d \leq y} |h_{1r}(d)| \ll \frac{xy^2}{(\log x)^3} \sum_{d=1}^{\infty} \frac{|h_{1r}(d)|}{d^2} \ll \frac{x}{\log x}.$$

Now using (29), for $0 < \alpha < 1$, we have

$$\begin{aligned} V_{12} &\ll \sum_{y < d \leq x+3} |h_{1r}(d)| \left(\frac{x}{d} + 1\right) \ll \sum_{y < d \leq x+3} |h_{1r}(d)| \ll (x+3) \sum_{d > y} \frac{|h_{1r}(d)|}{d} \\ &\ll \frac{(x+3)}{y^\alpha} \sum_{d=1}^{\infty} \frac{|h_{1r}(d)|}{d^{1-\alpha}} \ll \frac{x}{y^\alpha} \exp\left(c_2 \frac{r^\alpha}{\log r}\right) \end{aligned}$$

By taking $r = (\log \log x)^{\frac{1}{\alpha}}$, we have

$$V_{12} \ll xy^{-\alpha} y^{\frac{\alpha}{2}} = \frac{x}{(\log x)^{\frac{\alpha}{2}}}.$$

So as $x \rightarrow \infty$ we have,

$$V_1 = o(x).$$

From the similar calculation as Estimation of T_2 , we have

$$V_2 \ll (r \log r)^{-\frac{1}{2}} + \left(\sum_{r < p \leq x+3} \frac{|g_1(p) - 1|^2}{p} \right)^{1/2}.$$

From (12) and $r = (\log \log x)^{\frac{1}{\alpha}}$ we have as $x \rightarrow \infty$,

$$V_2 = o(x).$$

Which proves the required Theorem.

8. Proof of Applications

Proof of Application 1. We need the following lemma.

Lemma 8 ([3]). *Let $\{u_n\}_{n=1}^{\infty}$ and $\{v_n\}_{n=1}^{\infty}$ be two complex sequences such that*

$$\sum_{n=1}^{\infty} (|u_n|^2 + |v_n|) < \infty$$

Then,

$$\prod_{n=1}^{\infty} (1 + u_n + v_n) < \infty \text{ if and only if } \sum_{n=1}^{\infty} u_n < \infty.$$

Since $|g_j(p) - 1|^2 \leq 2(1 - \operatorname{Re}(g_j(p)))$, from (5) we have,

$$(31) \quad \sum_p \frac{|g_j(p) - 1|^2}{p} < \infty.$$

So by putting $r = \log x$, the error term in Theorem 1 is $o(1)$ as $x \rightarrow \infty$.
Now the infinite product $\prod_p w_p$ can be written as

$$\prod_p w_p = w_2 \prod_{p>2} w_p = w_2 \prod_p \left\{ 1 + \frac{g_1(p) + g_2(p) + g_3(p) - 3}{p} + O\left(\frac{1}{p^2}\right) \right\}.$$

From (5) and Lemma 8 we can say that, the above infinite product is convergent. Hence, by Theorem 1 Application is proved.

Proof of Application 2. We need the Lemma 8 and the following lemma:

Lemma 9 ([4], Lemma 6). *Let $F(n)$ be a polynomial as above of degree $v \geq 2$. we have the relation:*

$$\text{card} \left(n \leq x : F(n) \equiv 0 \pmod{p^{v-1}}, y_1 < p < \infty \right) = o(x),$$

when y_1 tends to infinity as $x \rightarrow \infty$.

Since $|g_j(p) - 1|^2 \leq 2(1 - \operatorname{Re}(g_j(p)))$, from (13) we have,

$$(32) \quad \sum_p \frac{|g_j(p) - 1|^2 \varrho_j(p)}{p} < \infty, j = 1, 2, 3.$$

From (13) and Lemma 8, we have $P_1(\gamma)$ and $P_2(\gamma)$ are convergent.

From Lemma 9, it is easy to see that as $p \rightarrow \infty$

$$(33) \quad (1 - g_j(p^{v_j-1})) \varrho_j(p^{v_j-1}) \rightarrow 0, j = 1, 2, 3.$$

So by putting $r = \log x$ and from (14), (32), (33), the error term in Theorem 2 is $o(1)$ as $x \rightarrow \infty$. Hence, by Theorem 2 Application is proved.

Proof of Application 3. It is easy to see that σ_a and ϕ are close to 1 for $a > 0$. The remainder term for both sums are estimate from the remainder term of Theorem 1 by choosing

$$\beta = 1 - \alpha = \min \left(\frac{1}{2k}, \frac{2 + c_{19}}{3} \right) \text{ and } r = c_{20}(\log x \log \log x)^{\frac{1}{\beta}},$$

for sufficiently small $c_{19}, c_{20} > 0$.

Hence, by Theorem 1 Application is proved.

Proof of Application 4 and 5. Since ϕ and σ_a are close to 1 for $a > 0$ then by applying Theorem 3 and Theorem 5 we get the Application 4 and Application 5 respectively.

Proof of Application 6. we need the following lemma:

Lemma 10. *Let p be an odd prime and $(a, p) = 1$, then $x^2 \equiv a \pmod{p^k}$ has exactly two solutions if a is a quadratic residue of p , and no solution if a is quadratic nonresidue of p . Further, if a is odd, then the congruence $x^2 \equiv a \pmod{2}$ is always solvable and has exactly one solution.*

We see that by Lemma 10, as $a > 0$, $p \rightarrow \infty$

$$(34) \quad \sum_p \frac{(g_j(p) - 1)\varrho_j(p)}{p} = \sum_p \frac{2}{p^{1+a}} < \infty, \quad (g_j(p) - 1)\varrho_j(p) = \frac{2}{p^a} \rightarrow 0,$$

where $g_j = \sigma_a$, $j = 1, 2, 3$, and

$$(35) \quad \sum_p \frac{(g_j(p) - 1)\varrho_j(p)}{p} = \sum_p \frac{-2}{p^2} < \infty, \quad (g_j(p) - 1)\varrho_j(p) = \frac{-2}{p} \rightarrow 0,$$

where $g_j = \phi$, $j = 1, 2, 3$.

The remainder term for both sum are estimate from the remainder term of Theorem 2 by choosing

$$\alpha = \frac{5 + c_{21}}{6} \quad \text{and} \quad r = c_{22}(\log x \log \log x)^{1/\alpha},$$

for sufficiently small $c_{21}, c_{22} > 0$.

Hence, by Theorem 2 Application is proved.

Proof of Application 7. we need the following lemma:

Lemma 11 ([3]). *Let $\{F_n\}_{n=1}^\infty$ be a sequence of distribution functions and $\{\phi_n\}_{n=1}^\infty$ then corresponding sequence of characteristic functions. Then F_n converges weakly to the distribution function F if and only if ϕ_n converges pointwise on $\mathbb{R}$ to a function ϕ which is continuous at 0. In addition, in this case, ϕ is the characteristic function of F and the convergence of ϕ_n to ϕ is uniform on any compact subset.*

The characteristic functions of the distribution (19) equal

$$(36) \quad \frac{1}{[x]} \sum_{n \leq x} \exp(it(f_1(n+3) + f_2(n+2) + f_3(n+1))).$$

Since

$$\sum_p \sum_{j=1}^3 \frac{\exp(itf_j(p)) - 1}{p} = t \sum_{j=1}^3 \sum_{|f_j(p)| \leq 1} \frac{f_j(p)}{p} + O\left(t^2 \sum_{j=1}^3 \left(\sum_{|f_j(p)| \leq 1} \frac{f_j^2(p)}{p} + \sum_{|f_j(p)| > 1} \frac{1}{p} \right)\right)$$

then from the convergence of the series (16), (17), (18) and from Lemma 8 we can say that the infinite product (20) converges for every t . This product is continuous at $t = 0$ because it converges uniformly for $|t| \leq T$ where $T > 0$ is arbitrary.

Since for $j = 1, 2, 3$

$$\sum_p \frac{|\exp(itf_j(p)) - 1|^2}{p} \ll t^2 \sum_{|f_j(p)| \leq 1} \frac{|f_j(p)|^2}{p} + \sum_{|f_j(p)| > 1} \frac{1}{p}$$

then from the convergence of (16) and (17) it follows that $S(r, x) \rightarrow 0$ when $r, x \rightarrow \infty$. Choosing $r = \log x$ in our Theorem 1 we get that the remainder term disappears when $x \rightarrow \infty$.

Thus the characteristic function (36) has the limit (20) for every real t and this limit is continuous at $t = 0$.

Therefore, by Lemma 11 Application is proved.

Proof of Application 8. We will use Lemma 8 and Lemma 11 to prove this application. The characteristic functions of the distribution (24) equal

$$(37) \quad \frac{1}{[x]} \sum_{n \leq x} \exp(it(f_1(F_1(n)) + f_2(F_2(n)) + f_3(F_3(n))))$$

Since

$$\begin{aligned} \sum_p \sum_{j=1}^3 \frac{(\exp(itf_j(p)) - 1) \varrho_j(p)}{p} &= t \sum_{j=1}^3 \sum_{|f_j(p)| \leq 1} \frac{f_j(p) \varrho_j(p)}{p} \\ &+ O\left(t^2 \sum_{j=1}^3 \sum_{|f_j(p)| \leq 1} \frac{f_j^2(p) \varrho_j(p)}{p}\right) + O\left(\sum_{j=1}^3 \sum_{|f_j(p)| > 1} \frac{\varrho_j(p)}{p}\right). \end{aligned}$$

then from the convergence of the series (21), (22), (23) and from Lemma 8 we can say that $P_1(\gamma)$ and $P_2(\gamma)$ are convergent for every real t . Further, the infinite product $P_1(\gamma)P_2(\gamma)$ is continuous at $t = 0$ because it converges uniformly for $|t| \leq T$ where $T > 0$ is arbitrary.

Since for $j = 1, 2, 3$

$$\sum_p \frac{|\exp(itf_j(p)) - 1|^2 \varrho_j(p)}{p} \ll t^2 \sum_{|f_j(p)| \leq 1} \frac{|f_j(p)|^2 \varrho_j(p)}{p} + \sum_{|f_j(p)| > 1} \frac{\varrho_j(p)}{p}$$

then from the convergence of (21) and (22) it follows that $S'(r, x), T(x) \rightarrow 0$ when $r, x \rightarrow \infty$.

Now from (24) and Lemma 9 it is easy to see that, $(\exp(itf_j(p^m)) - 1) \varrho_j(p^m) \rightarrow 0$ when $p \rightarrow \infty, m < v_j, j = 1, 2, 3$. Then $\frac{1}{x}C(r) \rightarrow 0$ as $r, x \rightarrow \infty$. Choosing $r = \log x$ in our Theorem 2 we get that the remainder term disappears when $x \rightarrow \infty$.

Thus the characteristic function (37) has the limit $P_1(\gamma)P_2(\gamma)$ for every real t and this limit is continuous at $t = 0$.

Therefore, by Lemma 11 Application is proved.

REFERENCES

- [1] P.D.T.A. Elliott *Probabilistic Number Theory I*, Springer, **239**, 1979.
- [2] P.D.T.A. Elliott *Extrapolating the mean-values of multiplicative functions*, Indagationes Mathematicae, **51**, 1989, pp. 409-420.
- [3] G. Tenenbaum *Introduction to Analytic and Probabilistic Number Theory* Cambridge University Press, 1995
- [4] I. Katái *On the distribution of Arithmetical functions*, Acta Mathematica Academiae Scientiarum Hungaricae, **20** (1-2), 1969, pp. 60-87.
- [5] H. Davenport *On some infinite series involving Arithmetical functions (II)*, The Quarterly Journal of Mathematics, **8**, 1937, pp. 313-320.
- [6] R. Warlimont *On multiplicative functions of absolute value ≤ 1* , Mathematische Nachrichten, **152**, 1991, pp. 113-120.
- [7] D. Wolke *Über die mittlere Verteilung der Werte zahlentheoretischer Funktionen auf Restklassen. I*, Mathematische Annalen, **202**, 1973, pp.1-25.

- [8] P. Erdős *On the sum $\sum_{k=1}^x d(f(k))$* , Journal of the London Mathematical Society, **s1-27**, 1952, pp. 7-15.
- [9] M. Tanaka *On the number of prime factors of integers I*, Japanese Journal of Mathematics, **25**, 1955, pp. 1-20.
- [10] G. Stepanauskas *The Mean values of Multiplicative Functions I*, Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae Sectio Mathematica, **18**, 1999, pp. 175-186.
- [11] G. Stepanauskas *The Mean values of Multiplicative Functions II*, Lithuanian Mathematical Journal, **37**, 1997, No. 4, pp. 443-451.
- [12] G. Stepanauskas *A note on the Liouville Function*, jános bolyai mathematical society, Budapest, 1999, pp. 245-248.
- [13] G. Stepanauskas *Mean values of Multiplicative Functions. III*, New Trends in Probability and Statistics, **4**, 1997, pp. 371-387.
- [14] Jonas Šiaulyš and G. Stepanauskas *On the Mean value of the product of Multiplicative functions with shifted argument*, Monatshefte für Mathematik, **150**, 2007, pp. 343-351.
- [15] Oleksiy Klurman *Correlations of Multiplicative functions and Applications*, arXiv.

INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI, CHENNAI 600113, INDIA
E-mail address, Pranendu Darbar: `dpranendu@imsc.res.in`